



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$8. \therefore \left[\frac{p}{v}(v+m)+n \right] \div (v-n) = \frac{p(v+m)+nv}{v(v-n)} = 20\frac{2}{3} \%.$$

Also solved by *M. E. GRABER*.

ALGEBRA.

125. Proposed by *LESLIE L. LOCKE*, Instructor in Mathematics, Michigan Agricultural College, Ingram County, Mich.

What special expedient will solve the system

$$\left. \begin{aligned} x^4 - y^4 &= 369 \dots (1) \\ x - y &= 1 \dots (2) \end{aligned} \right\} ?$$

Solution by *O. S. WESTCOTT*, Chicago, Ill.; *J. SCHEFFER*, A. M., Hagerstown, Md.; *J. K. ELWOOD*, A. M., Pittsburg, Pa.; *J. M. BOORMAN*, Woodmere, N. Y.; and the PROPOSER.

From (2), $y = x - 1$. Substituting this value of y in (1), $x^4 - (x-1)^4 = 369$ or $0 \cdot x^4 + 4x^3 - 6x^2 + 4x = 370$. Since the coefficient of x^4 is 0, therefore, if we regard the equation as of the fourth degree, one root is ∞ . Hence if $x = \infty$, $y = \infty$. The remaining three values of x are found from the equation $2x^3 - 3x^2 + 2x - 185 = 0$. This may be solved by the method of Tartaglia; or, by trial if 5 is substituted for x , the first member vanishes. Hence, $x = 5$. Dividing $2x^3 - 3x^2 + 2x - 185 = 0$, by $x - 5$, we have $2x^2 + 7x + 37 = 0$. $\therefore x = \frac{1}{2}[7 \pm \sqrt{(-247)}]$.

\therefore The values of x are ∞ , 5, and $\frac{1}{2}[7 \pm \sqrt{(-247)}]$, and the values of y are ∞ , 4, and $\frac{1}{2}[3 \pm \sqrt{(-247)}]$.

REMARK. The Proposer says his object in proposing this problem was, 1. To learn if there is a general method of solving such problems when factors can not be readily found, and, 2. To call attention to a fact that is not mentioned in many elementary text-books on Algebra, viz., the loss of a root by dividing one equation by another, or by subtracting one from another if thereby the degree of the equation is diminished.

The above contributors used some modifications in deriving the various steps in the solution of this problem, but these modifications were not considered of sufficient importance to warrant separate entries. Solutions were also received from *P. S. Berg*, *G. B. M. Zerr*, and *H. C. Whitaker*. Ed. F.

126. Proposed by *CHARLES C. CROSS*, Meredithville, Va.

A and B run a race; B, who runs slower than A by a miles in b hours, starts first by c minutes, and they get to the n -mile stone together. Required their rates of running. If $a=1$, $b=2$, $c=2$, and $n=4$, what is the result?

Solution by *GRANTLAND MURRAY*, Adjunct Professor of Mathematics, Emory College, Oxford, Ga.; *J. SCHEFFER*, A. M., Hagerstown, Md.; *H. C. WHITAKER*, A. M., Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pa.; *C. E. ARMENTROUT*, A. B., Professor of Latin and Mathematics, Rockingham Military Institute, Mt. Crawford, Va.; *P. S. BEEG*, Principal of Schools, Larimore, N. D.; *M. A. GRUBER*, A. M., War Department, Washington, D. C.; *C. ARTHUR LINDEMANN*, A. M., Professor of Mathematics and Science, Virginia Union University, Richmond, Va.; and *D. B. NORTHRUP*, Mandana, N. Y.

Let x = number of miles A runs per hour, then $x - a/b$ = number of miles B runs per hour.

n/x = number of hours A requires to run n miles, $n/(x - a/b)$ = number of B requires to run n miles.

$$\therefore \frac{n}{x - a/b} - \frac{n}{x} = \frac{c}{60}, \text{ or } bcx^2 - acx - 60an = 0; \text{ whence}$$

$$x = \frac{ac \pm \sqrt{(a^2c^2 + 240abcn)}}{2bc}.$$

Substituting the proposed values for a , b , c , and n gives, $x=8$, and $x-a/b=7\frac{1}{2}$.

Also solved by *G. B. M. ZERR*.

GEOMETRY.

153. Proposed by *WILLIAM HOOVER*, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

If P, P', Q, Q' be the extremities of two chords of a conic section, and both chords pass through the point A , show that the sum of the squares of the reciprocals of AP, AP', AQ, AQ' is constant.

No solution of this problem has been received.

156. Proposed by *F. M. McGAW*, A. M., Professor of Mathematics, Bordentown Military Institute, Bordentown, N. J.

To construct an equilateral triangle such that its vertices shall be in each of two parallel lines and a point fixed between these lines.

Solution by *G. I. HOPKINS*, A. M., Professor of Mathematics and Physics, High School, Manchester, N. H.

Let AB and CD be the two parallel lines, and F the fixed point between them. Through F draw HK perpendicular to CD . Make $\angle NMO=30^\circ$. Draw MN the perpendicular bisector of HK . Draw OS perpendicular to CD . Join F and P , and through P draw QR perpendicular to FP . Join QF and RF , then FQR is the required triangle.

PROOF. Triangles QOP and MFP are right triangles. $\angle QPO=\angle MPF$, being complements of the same $\angle QPM$.

\therefore these triangles are similar. $\therefore OP : MP :: QP : FP$, or by alternation $OP : QP :: MP : FP$. But these are homologous sides of the triangles OPM and QPF also.

\therefore these triangles are similar, since they are right triangles and the legs proportional. But the $\angle OMP$ is 30° and $\angle MOP$ is 60° .

$\therefore \angle QFP$ is 30° and $\angle FQP$ is 60° . Triangle FPR is easily shown to be equal to triangle FPQ .

$\therefore \angle FRP=60^\circ$. \therefore triangle FQR is equiangular and therefore equilateral.

Excellent solutions were received from *G. M. M. Zerr*, *H. C. Whitaker*, *J. Scheffer*, and *Theodore Linquist*. Professors Zerr's and Whitaker's solutions were by analytical geometry; Professor Scheffer's solution was by trigonometry and the application of algebra to geometry; and Professor Linquist, of the Kansas Agricultural College, gave a very good construction by pure geometry.

157. Proposed by *WILLIAM HOOVER*, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Find the locus of the center of a circle touching a given line and always passing through a given point.

